arXiv:1505.00668v2 [math.NT] 14 Jul 2015

Congruences for Catalan-Larcombe-French numbers

Xiao-Juan Ji[1] and Zhi-Hong Sun[2]

[1]School of Mathematical Sciences, Soochow University,
Suzhou, Jiangsu 215006, P.R. China
Email: Xiaojuanji2014@163.com

[2]School of Mathematical Sciences, Huaiyin Normal University,
Huaian, Jiangsu 223001, P.R. China
Email: zhihongsun@yahoo.com
Homepage: http://www.hytc.edu.cn/xsjl/szh

Abstract

Let $\{P_n\}$ be the Catalan-Larcombe-French numbers given by $P_0 = 1$, $P_1 = 8$ and $n^2P_n = 8(3n^2-3n+1)P_{n-1} - 128(n-1)^2P_{n-2}$ $(n \ge 2)$, and let $S_n = P_n/2^n$. In this paper we deduce congruences for $S_{mp^r} \pmod{p^{r+2}}$, $S_{mp^r-1} \pmod{p^r}$ and $S_{mp^r+1} \pmod{p^{2r}}$, where p is an odd prime and m, r are positive integers. We also prove that $S_{(p^2-1)/2} \equiv 0 \pmod{p^2}$ for any prime $p \equiv 5, 7 \pmod 8$, and show that $\{S_m\}$ is log-convex.

MSC: Primary 11A07, Secondary 05A10, 05A19
Keywords: congruence; Catalan-Larcombe-French number

1. Introduction

Let $\{P_n\}$ be the sequence given by

$$(1.1)\quad P_0 = 1,\ P_1 = 8 \quad \text{and} \quad (n+1)^2P_{n+1} = 8(3n^2+3n+1)P_n - 128n^2P_{n-1}\ (n \ge 1).$$

The numbers P_n are called Catalan-Larcombe-French numbers since Catalan first defined P_n in [C], and in [LF1] Larcombe and French proved that

$$(1.2)\qquad P_n = 2^n \sum_{k=0}^{[n/2]} (-4)^k \binom{2n-2k}{n-k}^2 \binom{n-k}{k} = \sum_{k=0}^{n} \frac{\binom{2k}{k}^2\binom{2n-2k}{n-k}^2}{\binom{n}{k}},$$

where $[x]$ is the greatest integer not exceeding x. The numbers P_n occur in the theory of elliptic integrals, and are related to the arithmetic-geometric-mean. See [LF1] and A053175 in Sloane's database "The On-Line Encyclopedia of Integer Sequences".

Let $\{f_n\}$ be the Franel numbers given by $f_n = \sum_{k=0}^{n} \binom{n}{k}^3$ $(n = 0, 1, 2, \dots)$, and let $\{S_n\}$ be defined by

$$(1.3)\quad S_0 = 1,\ S_1 = 4 \quad \text{and} \quad (n+1)^2S_{n+1} = 4(3n^2+3n+1)S_n - 32n^2S_{n-1}\ (n \ge 1).$$

Comparing (1.3) with (1.1), we see that

$$S_n = \frac{P_n}{2^n}. \tag{1.4}$$

Zagier noted that

$$S_n = \sum_{k=0}^{[n/2]} \binom{2k}{k}^2 \binom{n}{2k} 4^{n-2k}. \tag{1.5}$$

In this paper we investigate the properties of S_n instead of P_n since S_n is an Apépy-like sequence. As observed by V. Jovovic in 2003 (see [LF2]),

$$S_n = \sum_{k=0}^{n} \binom{n}{k}\binom{2k}{k}\binom{2n-2k}{n-k} \quad (n = 0, 1, 2, \ldots). \tag{1.6}$$

Recently Z.W. Sun stated that

$$S_n = \sum_{k=0}^{n} \binom{2k}{k}^2 \binom{k}{n-k}(-4)^{n-k} = \frac{1}{(-2)^n}\sum_{k=0}^{n} \binom{2k}{k}\binom{2n-2k}{n-k}\binom{k}{n-k}(-4)^k. \tag{1.7}$$

The first few values of S_n are shown below:

$$S_0 = 1,\ S_1 = 4,\ S_2 = 20,\ S_3 = 112,\ S_4 = 676,\ S_5 = 4304,$$
$$S_6 = 28496,\ S_7 = 194240,\ S_8 = 1353508,\ S_9 = 9593104.$$

Let p be an odd prime. In [JLF], Jarvis, Larcombe and French proved that if $n = a_rp^r + \cdots + a_1p + a_0$ with $a_0, a_1, \ldots, a_r \in \{0, 1, \ldots, p-1\}$, then

$$P_n \equiv P_{a_r} \cdots P_{a_1} P_{a_0} \pmod p. \tag{1.8}$$

In [JV] Jarvis and Verrill showed that

$$P_n \equiv (-1)^{\frac{p-1}{2}} 128^n P_{p-1-n} \pmod p \quad \text{for} \quad n = 0, 1, \ldots, p-1 \tag{1.9}$$

and

$$P_{mp^r} \equiv P_{mp^{r-1}} \pmod{p^r} \quad \text{for} \quad m, r \in \mathbb{Z}^+, \tag{1.10}$$

where $\mathbb{Z}^+$ is the set of positive integers. In [OS] Osburn and Sahu stated that

$$S_{mp^r} \equiv S_{mp^{r-1}} \pmod{p^{2r}} \quad \text{for} \quad m, r \in \mathbb{Z}^+. \tag{1.11}$$

But they did not give the details for the proof. In this paper we will prove (1.11) in an elementary and natural manner. Let $\varphi(n)$ be the Euler's totient function. Since $P_n = 2^nS_n$, from (1.11) we deduce a congruence for $P_{mp^r} - 2^{m\varphi(p^r)}P_{mp^{r-1}} \pmod{p^{2r}}$, which improves (1.10). Thus (1.11) is a vast generalization of (1.10).

In [S3] the second author established some identities involving S_n. For example,

$$\sum_{k=0}^{n} \binom{n}{k}(-1)^k \frac{S_k}{8^k} = \frac{S_n}{8^n} \quad \text{and} \quad \sum_{k=0}^{2n} \binom{2n}{k}\binom{2n+k}{k}(-8)^{2n-k} S_k = (-1)^n \binom{2n}{n}^3. \tag{1.12}$$

For a prime p let $\mathbb{Z}_p$ denote the set of those rational numbers whose denominator is not divisible by p. Let p be an odd prime, $n \in \mathbb{Z}_p$ and $n \not\equiv 0, -16 \pmod p$. In [S3] the second author proved that

$$\sum_{k=0}^{p-1} \binom{2k}{k} \frac{S_k}{(n+16)^k} \equiv \Big(\frac{n(n+16)}{p}\Big) \sum_{k=0}^{p-1} \frac{\binom{2k}{k}^2\binom{4k}{2k}}{n^{2k}} \pmod p, \tag{1.13}$$

where $(\frac{a}{p})$ is the Legendre symbol.

Let $r \in \mathbb{Z}^+$ and p be a prime with $p \equiv 5, 7 \pmod 8$. In [S3] the second author conjectured that

$$S_{\frac{p^r-1}{2}} \equiv 0 \pmod {p^r} \quad \text{and} \quad f_{\frac{p^r-1}{2}} \equiv 0 \pmod {p^r}. \tag{1.14}$$

In this paper we prove (1.14) in the case $r = 2$.

Let $\{E_n\}$ be the Euler numbers given by

$$E_{2n-1} = 0,\ E_0 = 1 \quad \text{and} \quad \sum_{k=0}^{n} \binom{2n}{2k} E_{2k} = 0\ (n \ge 1).$$

Suppose that p is an odd prime and $n \in \mathbb{Z}^+$. In this paper we determine $S_{mp^r} \pmod {p^{r+2}}$ by showing that

$$S_{np} - S_n \equiv \begin{cases} 8n^2 S_{n-1}(-1)^{\frac{p-1}{2}} p^2 E_{p-3} \pmod {p^3} & \text{if } p > 3 \text{ and } p \nmid n, \\ 9(n-1)S_n \pmod {p^3} & \text{if } p = 3 \text{ and } 3 \nmid n, \\ 0 \pmod {p^{3+\text{ord}_p n}} & \text{if } p \mid n, \end{cases} \tag{1.15}$$

where $\text{ord}_p n$ is the unique nonnegative integer α such that $p^\alpha \mid n$ and $p^{\alpha+1} \nmid n$. We also show that

$$S_{mp^r+1} \equiv 4(mp^r+1)S_{mp^r-1} \pmod {p^{2r}} \text{ and } S_{mp^r-1} \equiv (-1)^{\frac{p-1}{2}} S_{mp^{r-1}-1} \pmod {p^r}. \tag{1.16}$$

In Section 4 we prove the second author's conjecture (see [S3])

$$S_m^2 < S_{m+1}S_{m-1} < \big(1 + \frac{1}{m(m-1)}\big) S_m^2 \quad \text{for} \quad m = 2, 3, \ldots.$$

2. Basic lemmas

Lemma 2.1 (Lucas theorem [M]). *Let p be an odd prime. Suppose $a = a_r p^r + \cdots + a_1 p + a_0$ and $b = b_r p^r + \cdots + b_1 p + b_0$, where $a_r, \ldots, a_0, b_r, \ldots, b_0 \in \{0, 1, \ldots, p-1\}$. Then*

$$\binom{a}{b} \equiv \binom{a_r}{b_r} \cdots \binom{a_0}{b_0} \pmod p.$$

Lemma 2.2. *Let p be an odd prime and $a, b \in \mathbb{Z}^+$. Suppose $a_0, b_0 \in \{0, 1, \ldots, p-1\}$. Then*

$$\binom{ap+a_0}{bp+b_0} \equiv \binom{a}{b}\binom{a_0}{b_0} \pmod p.$$

Proof. Assume $a = a_rp^r + \cdots + a_2p + a_1$ and $b = b_rp^r + \cdots + b_2p + b_1$, where $a_r, \ldots, a_1, b_r, \ldots, b_1 \in \{0, 1, \ldots, p-1\}$. By Lucas theorem,

$$\binom{ap + a_0}{bp + b_0} \equiv \binom{a_r}{b_r} \cdots \binom{a_1}{b_1}\binom{a_0}{b_0} \equiv \binom{a}{b}\binom{a_0}{b_0} \pmod p.$$

This is the result.

Lemma 2.3 (Kazandzidis' congruence [M]). *Let $p \geq 5$ be a prime and $m, n \in \mathbb{Z}^+$. Then*

$$\binom{mp}{np} \equiv \binom{m}{n} \pmod{p^3}.$$

Lemma 2.4 ([Su, Lemma 2.1]). *Let p be an odd prime and $k \in \{1, 2, \ldots, p-1\}$. Then*

$$\binom{2k}{k}\binom{2(p-k)}{p-k} \equiv \begin{cases} -\frac{2p}{k} \pmod{p^2} & \text{if } k < \frac{p}{2}, \\ \frac{2p}{k} \pmod{p^2} & \text{if } k > \frac{p}{2}. \end{cases}$$

Let $\{B_n\}$ be the Bernoulli numbers defined by $B_0 = 1$ and $\sum_{k=0}^{n-1}\binom{n}{k}B_k = 0$ $(n \geq 2)$. It is known that $B_{2k+1} = 0$ for $k \in \mathbb{Z}^+$. For $m, n \in \mathbb{Z}^+$ it is well known that

$$\sum_{k=0}^{n-1} k^m = \frac{1}{m+1}\sum_{k=1}^{m+1}\binom{m+1}{k}B_k n^{m+1-k}. \tag{2.1}$$

By the Staudt-Clausen theorem, $B_{2k} \in \mathbb{Z}_p$ for $2k \not\equiv 0 \pmod{p-1}$, and $pB_{2k} \in \mathbb{Z}_p$ for $2k \equiv 0 \pmod{p-1}$. See [MOS].

Let $\{E_n(x)\}$ be the Euler polynomials given by

$$E_n(x) = \frac{1}{2^n}\sum_{k=0}^{n}\binom{n}{k}(2x-1)^{n-k}E_k.$$

Then $E_n = 2^nE_n(\frac{1}{2})$. It is known that (see [MOS])

$$\sum_{k=0}^{n-1}(-1)^k k^m = \frac{E_m(0) - (-1)^nE_m(n)}{2}.$$

Lemma 2.5 ([S2, Lemma 2.2]). *Let p be an odd prime, $a \in \mathbb{Z}_p$, $a \not\equiv 0 \pmod p$ and $k \in \{1, 2, \ldots, p-2\}$. Then*

$$\sum_{r=1}^{\langle a\rangle_p}\frac{(-1)^r}{r^k} \equiv -\frac{(2^{p-k}-1)B_{p-k}}{p-k} + \frac{1}{2}(-1)^{\langle a\rangle_p + k}E_{p-1-k}(-a) \pmod p,$$

where $\langle a\rangle_p \in \{0, 1, \ldots, p-1\}$ is given by $a \equiv \langle a\rangle_p \pmod p$.

Lemma 2.6. *Let p be an odd prime, $k, m \in \mathbb{Z}^+$ and $\frac{p}{2} < k < p$. Then*

$$\binom{2mp + 2k}{mp + k} \equiv (2m+1)\binom{2m}{m}\binom{2k}{k} \pmod{p^2}.$$

Proof. Clearly

$$\begin{aligned}
&\frac{(2m+1)p((2m+1)p-1)\cdots((m+1)p+1)(m+1)p}{(mp)!}\\
&=\frac{((2m+1)p)(2mp)\cdots((m+1)p)}{p\cdot(2p)\cdots(mp)}\cdot\frac{\prod_{r=m+1}^{2m}(rp+1)\cdots(rp+p-1)}{\prod_{r=0}^{m-1}(rp+1)\cdots(rp+p-1)}\\
&=p(2m+1)\binom{2m}{m}\cdot\frac{\prod_{r=m+1}^{2m}(rp+1)\cdots(rp+p-1)}{\prod_{r=0}^{m-1}(rp+1)\cdots(rp+p-1)}\\
&\equiv p(2m+1)\binom{2m}{m}\frac{(p-1)!^m}{(p-1)!^m}=p(2m+1)\binom{2m}{m}\pmod{p^2}.
\end{aligned}$$

Thus

$$\begin{aligned}
&\binom{2mp+2k}{mp+k}\\
&=\frac{(2m+1)p((2m+1)p-1)\cdots((m+1)p+1)(m+1)p}{(mp)!}\\
&\quad\times\frac{(2mp+2k)\cdots(2mp+p+1)((m+1)p-1)\cdots((m+1)p-(p-1-k))}{(mp+1)\cdots(mp+k)}\\
&\equiv p(2m+1)\binom{2m}{m}\frac{(2k)(2k-1)\cdots(p+1)(p-1)(p-2)\cdots(k+1)}{k!}\\
&=(2m+1)\binom{2m}{m}\binom{2k}{k}\pmod{p^2}.
\end{aligned}$$

This proves the lemma.

Lemma 2.7. *For any positive integer n we have*

$$S_n=2\sum_{k=1}^{n}\binom{n-1}{k-1}\binom{2k}{k}\binom{2n-2k}{n-k}.$$

Proof. Since

$$\begin{aligned}
\sum_{k=0}^{n}(2k-n)\binom{n}{k}\binom{2k}{k}\binom{2n-2k}{n-k}&=\sum_{k=0}^{n}(2(n-k)-n)\binom{n}{k}\binom{2k}{k}\binom{2n-2k}{n-k}\\
&=-\sum_{k=0}^{n}(2k-n)\binom{n}{k}\binom{2k}{k}\binom{2n-2k}{n-k},
\end{aligned}$$

we see that

$$\sum_{k=0}^{n}(2k-n)\binom{n}{k}\binom{2k}{k}\binom{2n-2k}{n-k}=0 \tag{2.2}$$

and so

$$nS_n=\sum_{k=0}^{n}2k\binom{n}{k}\binom{2k}{k}\binom{2n-2k}{n-k}=2n\sum_{k=1}^{n}\binom{n-1}{k-1}\binom{2k}{k}\binom{2n-2k}{n-k}.$$

This yields the result.

Lemma 2.8. *Let* $m \in \mathbb{Z}$ *and* $k, n, p \in \mathbb{Z}^+$. *Then*

$$\binom{mp^r - 1}{k} = (-1)^{k-[\frac{k}{p}]} \binom{mp^{r-1} - 1}{[k/p]} \prod_{\substack{i=1\\ p\nmid i}}^{k} \Big(1 - \frac{mp^r}{i}\Big)$$

and so

$$\binom{mp^r}{np} = (-1)^{n(p-1)} \binom{mp^{r-1}}{n} \prod_{\substack{i=1\\ p\nmid i}}^{np-1} \Big(1 - \frac{mp^r}{i}\Big).$$

Proof. Clearly,

$$\begin{aligned}\binom{mp^r - 1}{k} &= \prod_{i=1}^{k} \frac{mp^r - i}{i} = \prod_{\substack{i=1\\ p\nmid i}}^{k} \frac{mp^r - i}{i} \prod_{i=1}^{[k/p]} \frac{mp^r - pi}{pi} \\ &= \prod_{\substack{i=1\\ p\nmid i}}^{k} \frac{mp^r - i}{i} \prod_{i=1}^{[k/p]} \frac{mp^{r-1} - i}{i} \\ &= (-1)^{k-[\frac{k}{p}]} \prod_{\substack{i=1\\ p\nmid i}}^{k} \Big(1 - \frac{mp^r}{i}\Big) \cdot \binom{mp^{r-1} - 1}{[k/p]}.\end{aligned}$$

Taking $k = np - 1$ in the above we see that

$$\binom{mp^r}{np} = \frac{mp^r}{np} \binom{mp^r - 1}{np - 1} = \frac{mp^{r-1}}{n} \binom{mp^{r-1} - 1}{n - 1} \cdot (-1)^{np-1-(n-1)} \prod_{\substack{i=1\\ p\nmid i}}^{np-1} \Big(1 - \frac{mp^r}{i}\Big).$$

This yields the remaining part.

Lemma 2.9. *Let* $m, n \in \mathbb{Z}^+$.*Then*

$$\binom{3m}{3n} \equiv \binom{m}{n} (1 + 9mn^2 - 9m^2 n) \pmod{27}.$$

Proof. By Lemma 2.8,

$$\begin{aligned}\binom{3m}{3n} &= \binom{m}{n} \prod_{\substack{k=1\\ 3\nmid k}}^{3n-1} (1 - \frac{3m}{k}) \\ &\equiv \binom{m}{n} \Big(1 - 3m \sum_{\substack{k=1\\ 3\nmid k}}^{3n-1} \frac{1}{k} + 9m^2 \sum_{\substack{1\le i<j\le 3n-1\\ 3\nmid ij}} \frac{1}{ij}\Big) \pmod{27}.\end{aligned}$$

Clearly

$$\begin{aligned}2 \sum_{\substack{1\le i<j\le 3n-1\\ 3\nmid ij}} \frac{1}{ij} &= \Big(\sum_{\substack{i=1\\ 3\nmid i}}^{3n-1} \frac{1}{i}\Big)^2 - \sum_{\substack{i=1\\ 3\nmid i}}^{3n-1} \frac{1}{i^2} \equiv \Big(\sum_{i=1}^{3n-1} i\Big)^2 - \sum_{\substack{i=1\\ 3\nmid i}}^{3n-1} 1 \\ &= \Big(\frac{(3n-1)3n}{2}\Big)^2 - (3n - 1 - (n-1)) \equiv -2n \pmod{3}.\end{aligned}$$

Thus,

$$\binom{3m}{3n} \equiv \binom{m}{n}\Big(1 - 3m\sum_{\substack{k=1\\3\nmid k}}^{3n-1}\frac{1}{k} - 9m^2n\Big) \pmod{27}.$$

By Euler's Theorem, for $k \not\equiv 0 \pmod 3$ we have $k^6 = k^{\varphi(9)} \equiv 1 \pmod 9$. Thus

$$\begin{aligned}\sum_{\substack{k=1\\3\nmid k}}^{3n-1}\frac{1}{k} &\equiv \sum_{k=1}^{3n-1}k^5 = \frac{B_6(3n)-B_6}{6} = \frac{1}{6}\sum_{k=1}^{6}\binom{6}{k}(3n)^kB_{6-k}\\ &\equiv \frac{1}{6}\binom{6}{2}(3n)^2B_4 = -\frac{3}{4}n^2 \equiv -3n^2 \pmod 9.\end{aligned}$$

Hence

$$\begin{aligned}\binom{3m}{3n} &\equiv \binom{m}{n}(1+9mn^2-9m^2n)\\ &\equiv \begin{cases}\binom{m}{n} \pmod{27} & \text{if } 3 \mid mn(m-n)\ ,\\ \binom{m}{n}(1+9n) \pmod{27} & \text{if } 3\nmid mn \text{ and } 3 \mid m+n.\end{cases}\end{aligned}$$

Lemma 2.10. *Let p be an odd prime, $r, m \in \mathbb{Z}^+$ and $s \in \{0, 1, \dots, mp^{r-1}-1\}$. Then*

$$\binom{mp^r}{sp} \equiv \binom{mp^{r-1}}{s} \pmod{p^{2r}}.$$

Proof. By Lemma 2.8,

$$\binom{mp^r}{sp} = \binom{mp^{r-1}}{s}\prod_{\substack{i=1\\p\nmid i}}^{sp-1}\Big(1-\frac{mp^r}{i}\Big) \equiv \binom{mp^{r-1}}{s}\Big(1 - mp^r\sum_{\substack{i=1\\p\nmid i}}^{sp-1}\frac{1}{i}\Big) \pmod{p^{2r}}.$$

Set $l = \mathrm{ord}_p s$ and $s = p^l s_0$. Since $B_1 = -\frac{1}{2}$, $B_{2s+1} = 0$ $(s \geq 1)$, $pB_k \in \mathbb{Z}_p$ and $\varphi(p^{l+1}) \geq l+2$ we see that

$$\begin{aligned}\sum_{\substack{i=1\\p\nmid i}}^{sp-1}\frac{1}{i} &\equiv \sum_{\substack{i=1\\p\nmid i}}^{sp-1} i^{\varphi(p^{l+1})-1} \equiv \sum_{i=1}^{sp-1} i^{\varphi(p^{l+1})-1}\\ &= \frac{1}{\varphi(p^{l+1})}\sum_{j=1}^{\varphi(p^{l+1})}\binom{\varphi(p^{l+1})}{j}(s_0p^{l+1})^jB_{\varphi(p^{l+1})-j}\\ &= (s_0p^{l+1})^{\varphi(p^{l+1})-1}B_1 + \frac{1}{\varphi(p^{l+1})}\sum_{j=1}^{\varphi(p^{l+1})/2}\binom{\varphi(p^{l+1})}{2j}(s_0p^{l+1})^{2j}B_{\varphi(p^{l+1})-2j}\\ &\equiv \frac{s_0}{p-1}\sum_{j=1}^{\varphi(p^{l+1})/2}\binom{\varphi(p^{l+1})}{2j}(s_0p^{l+1})^{2j-1}\cdot pB_{\varphi(p^{l+1})-2j} \equiv 0 \pmod{p^{l+1}}.\end{aligned} \tag{2.3}$$

If $l \ge r-1$, then $r+l+1 \ge 2r$ and so

$$\binom{mp^r}{sp} \equiv \binom{mp^{r-1}}{s}\Big(1 - mp^r \sum_{\substack{i=1\\ p\nmid i}}^{sp-1}\frac{1}{i}\Big) \equiv \binom{mp^{r-1}}{s} \pmod{p^{2r}}.$$

If $0 \le l < r-1$, then $\binom{mp^{r-1}}{s} = \frac{mp^{r-1}}{s_0 p^l}\binom{mp^{r-1}-1}{s-1} \equiv 0 \pmod{p^{r-1-l}}$ and so

$$\binom{mp^r}{sp} \equiv \binom{mp^{r-1}}{s} - mp^r \binom{mp^{r-1}}{s}\sum_{\substack{i=1\\ p\nmid i}}^{sp-1}\frac{1}{i} \equiv \binom{mp^{r-1}}{s} \pmod{p^{2r}}.$$

This completes the proof.

Lemma 2.11. *Let p be an odd prime, $r, m \in \mathbb{Z}^+$ and $s \in \{1, 2, \dots, mp^{r-1}\}$. Then*

$$\begin{aligned}&\binom{mp^{r-1}}{s}\binom{2sp}{sp}\binom{2(mp^{r-1}-s)p}{(mp^{r-1}-s)p}\\ &\equiv \begin{cases}\binom{m}{s}\binom{2s}{s}\binom{2(m-s)}{m-s}(1+9m) \pmod{p^{r+2}} & \text{if } r=1 \text{ and } p=3,\\ \binom{mp^{r-1}}{s}\binom{2s}{s}\binom{2(mp^{r-1}-s)}{mp^{r-1}-s} \pmod{p^{r+2}} & \text{if } r>1 \text{ or } p>3.\end{cases}\end{aligned}$$

Proof. For $r=1$ the result follows from Lemmas 2.3 and 2.9. Now assume $r \ge 2$. If $p \nmid s$, then $\binom{mp^{r-1}}{s} = \frac{mp^{r-1}}{s}\binom{mp^{r-1}-1}{s-1} \equiv 0 \pmod{p^{r-1}}$. By Lemmas 2.3 and 2.9,

$$\binom{2sp}{sp}\binom{2(mp^{r-1}-s)p}{(mp^{r-1}-s)p} \equiv \binom{2s}{s}\binom{2(mp^{r-1}-s)}{mp^{r-1}-s} \pmod{p^3}.$$

Thus the result is true. Now assume that $p \mid s$, $l = \mathrm{ord}_p s$ and $s = p^l s_0$. For $1 \le l < r-1$, using Lemma 2.10 we see that

$$\binom{2sp}{sp}\binom{2(mp^{r-1}-s)p}{(mp^{r-1}-s)p} \equiv \binom{2s}{s}\binom{2(mp^{r-1}-s)}{mp^{r-1}-s} \pmod{p^{2l+2}}.$$

Since $\binom{mp^{r-1}}{s} = \frac{mp^{r-1}}{p^l s_0}\binom{mp^{r-1}-1}{s-1} \equiv 0 \pmod{p^{r-1-l}}$ and $r-1-l+2l+2 = r+l+1 \ge r+2$, the result is true in this case. For $l \ge r$ we see that $p^r \mid sp$ and $p^r \mid (mp^{r-1}-s)p$. Thus applying Lemma 2.10 we deduce that

$$\binom{2sp}{sp}\binom{2(mp^{r-1}-s)p}{(mp^{r-1}-s)p} \equiv \binom{2s}{s}\binom{2(mp^{r-1}-s)}{mp^{r-1}-s} \pmod{p^{2r}}.$$

As $2r \ge r+2$, the result is also true. The proof is now complete.

Lemma 2.12. *Let p be an odd prime, $m, r \in \mathbb{Z}^+$ and $k \in \{0, \dots, mp^r\}$. Then*

$$k\binom{2k}{k}\binom{2(mp^r-k)}{mp^r-k} \equiv 0 \pmod{p^r}.$$

Proof. Suppose $s = [\frac{k}{p}]$ and $t = k - sp$. Then $t \in \{0, 1, \dots, p-1\}$. We first assume $p \nmid k$. That is, $t > 0$. Let us consider the case $r = 1$. By Lemma 2.2, for $1 \le t < \frac{p}{2}$,

$$\binom{2k}{k}\binom{2(mp-k)}{mp-k} = \binom{2k}{k}\binom{(2(m-s)-1)p+p-2t}{(m-s-1)p+p-t}$$

$$\equiv \binom{2k}{k}\binom{2(m-s)-1}{m-s-1}\binom{p-2t}{p-t} = 0 \pmod{p},$$

and for $t > \frac{p}{2}$,

$$\begin{aligned}\binom{2k}{k}\binom{2(mp-k)}{mp-k} &= \binom{(2s+1)p+2t-p}{sp+t}\binom{2(mp-k)}{mp-k}\\ &\equiv \binom{2s+1}{s}\binom{2t-p}{t}\binom{2(mp-k)}{mp-k} = 0 \pmod{p}.\end{aligned}$$

Thus the result is true for $r = 1$.

Now assume $p \nmid k$ and $r \geq 2$. Suppose that for $n < r$ and $k \in \{1, 2, \ldots, mp^n - 1\}$ we have

$$\binom{2k}{k}\binom{2(mp^n-k)}{mp^n-k} \equiv 0 \pmod{p^n}.$$

When $p \mid s$, by the inductive hypothesis we have

$$\begin{aligned}&\binom{2s}{s}\binom{2mp^{r-1}-2s-2}{mp^{r-1}-s-1}(2mp^{r-1}-2s-1)p\\ &= \frac{s+1}{2(2s+1)}(2mp^{r-1}-2s-1)p\binom{2s+2}{s+1}\binom{2mp^{r-1}-2s-2}{mp^{r-1}-s-1} \equiv 0 \pmod{p^r}.\end{aligned}$$

When $p \nmid s$, by the inductive hypothesis we obtain

$$\begin{aligned}&\binom{2s}{s}\binom{2mp^{r-1}-2s-2}{mp^{r-1}-s-1}(2mp^{r-1}-2s-1)p\\ &= \frac{mp^{r-1}-s}{2}p\binom{2s}{s}\binom{2mp^{r-1}-2s}{mp^{r-1}-s} \equiv 0 \pmod{p^r}.\end{aligned}$$

Suppose $k \in \{1, 2, \ldots, mp^r - 1\}$. For $t < \frac{p}{2}$, from the above we see that

$$\begin{aligned}\binom{2k}{k}\binom{2(mp^r-k)}{mp^r-k} &= \binom{2sp+2t}{sp+t}\binom{(2mp^{r-1}-2s-1)p+p-2t}{(mp^{r-1}-s-1)p+p-t}\\ &= \binom{2s}{s}\binom{2mp^{r-1}-2s-2}{mp^{r-1}-s-1}(2mp^{r-1}-2s-1)pQ \equiv 0 \pmod{p^r},\end{aligned}$$

and for $t > \frac{p}{2}$,

$$\begin{aligned}\binom{2k}{k}\binom{2(mp^r-k)}{mp^r-k} &= \binom{(2s+1)p+2t-p}{sp+t}\binom{(2mp^{r-1}-2s-2)p+2p-2t}{(mp^{r-1}-s-1)p+p-t}\\ &= Q\binom{2s}{s}\binom{2mp^{r-1}-2s-2}{mp^{r-1}-s-1}(2s+1)p\\ &\equiv -Q\binom{2s}{s}\binom{2mp^{r-1}-2s-2}{mp^{r-1}-s-1}(2mp^{r-1}-2s-1)p\\ &\equiv 0 \pmod{p^r},\end{aligned}$$

where

$$Q = \frac{\sum\limits_{\substack{i=1\\ p\nmid i}}^{2sp+2t} i \sum\limits_{\substack{i=1\\ p\nmid i}}^{2(mp^r-sp-t)} i}{\sum\limits_{\substack{i=1\\ p\nmid i}}^{sp+t} i^2 \sum\limits_{\substack{i=1\\ p\nmid i}}^{mp^r-sp-t} i^2} \in \mathbb{Z}_p.$$

Hence the result is true for $n = r$. Summarizing the above we prove the result in the case $p \nmid k$.

Now we assume $p \mid k$. Set $l = \mathrm{ord}_p k$ and $k = p^l k_0$. Then $k_0 \in \{1, \ldots, mp^{r-l} - 1\}$ and $p \nmid k_0$. For $l \ge r$ obviously we have $k\binom{2k}{k}\binom{2(mp^r-k)}{mp^r-k} \equiv 0 \pmod{p^r}$. For $1 \le l \le r-1$, since $p \nmid k_0$, from the above we deduce that

$$k\binom{2k}{k}\binom{2(mp^r-k)}{mp^r-k} = Wp^l\binom{2k_0}{k_0}\binom{2mp^{r-l}-2k_0}{mp^{r-l}-k_0} \equiv 0 \pmod{p^r},$$

where $W \in \mathbb{Z}_p$. The proof is now complete.

Lemma 2.13. *Let p be an odd prime, $r, m \in \mathbb{Z}^+$ and $s \in \{0, 1, \ldots, mp^{r-1} - 1\}$. Then*

$$\binom{2sp+p-1}{sp+\frac{p-1}{2}}\binom{2(mp^{r-1}-s-1)p+p-1}{(mp^{r-1}-s-1)p+\frac{p-1}{2}} \equiv \binom{2s}{s}\binom{2(mp^{r-1}-s-1)}{mp^{r-1}-s-1} \pmod{p^r}.$$

Proof. Using Lemma 2.8 and the identity

$$\binom{a-b}{c-d}\binom{b}{d} = \frac{\binom{a}{c}\binom{c}{d}\binom{a-c}{b-d}}{\binom{a}{b}} \tag{2.4}$$

we have

$$\begin{aligned}
&\binom{2sp+p-1}{sp+\frac{p-1}{2}}\binom{2(mp^{r-1}-s-1)p+p-1}{(mp^{r-1}-s-1)p+(p-1)/2}\\
&= \frac{mp^{r-1}}{2(2s+1)} \cdot \frac{\binom{2mp^r}{mp^r}\binom{mp^r-1}{sp+(p-1)/2}^2}{\binom{2mp^r-1}{2sp+p}}\\
&= \frac{mp^{r-1}}{2(2s+1)} \cdot \frac{\binom{2mp^{r-1}}{mp^{r-1}}\binom{mp^{r-1}-1}{s}^2}{\binom{2mp^{r-1}-1}{2s+1}} \cdot \frac{\prod\limits_{\substack{i=1\\ p\nmid i}}^{mp^r} \frac{2mp^r-i}{i} \Big(\prod\limits_{\substack{i=1\\ p\nmid i}}^{sp+(p-1)/2} \frac{mp^r-i}{i}\Big)^2}{\prod\limits_{\substack{i=1\\ p\nmid i}}^{2sp+p} \frac{2mp^r-i}{i}},
\end{aligned}$$

and

$$\binom{2s}{s}\binom{2(mp^{r-1}-s-1)}{mp^{r-1}-s-1} = \frac{mp^{r-1}}{2(2s+1)} \cdot \frac{\binom{2mp^{r-1}}{mp^{r-1}}\binom{mp^{r-1}-1}{s}^2}{\binom{2mp^{r-1}-1}{2s+1}}.$$

Since

$$\frac{\binom{2sp+p-1}{sp+(p-1)/2}\binom{2(mp^{r-1}-s-1)p+p-1}{(mp^{r-1}-s-1)p+(p-1)/2}}{\binom{2s}{s}\binom{2(mp^{r-1}-s-1)}{mp^{r-1}-s-1}} = \frac{\prod\limits_{\substack{i=1\\ p\nmid i}}^{mp^r} \frac{2mp^r-i}{i} \prod\limits_{\substack{i=1\\ p\nmid i}}^{sp+(p-1)/2} (\frac{mp^r-i}{i})^2}{\prod\limits_{\substack{i=1\\ p\nmid i}}^{2sp+p} \frac{2mp^r-i}{i}} \equiv \frac{\prod\limits_{\substack{i=1\\ p\nmid i}}^{mp^r}(-1) \prod\limits_{\substack{i=1\\ p\nmid i}}^{sp+(p-1)/2} (-1)^2}{\prod\limits_{\substack{i=1\\ p\nmid i}}^{2sp+p}(-1)}$$
$$= 1 \pmod{p^r},$$

we obtain the result.

Lemma 2.14. *Let p be an odd prime, $r, m \in \mathbb{Z}^+$ and $s \in \{0, 1, \ldots, mp^{r-1}-1\}$. Then*

$$\binom{mp^r}{sp}\binom{2sp}{sp}\binom{2mp^r-2sp}{mp^r-sp} \equiv \binom{mp^{r-1}}{s}\binom{2s}{s}\binom{2mp^{r-1}-2s}{mp^{r-1}-s} \pmod{p^{2r}}.$$

Proof. Set $l = \mathrm{ord}_p s$ and $s = p^l s_0$. By (2.3), (2.4), Lemmas 2.8 and 2.10, we have

$$\frac{\binom{2sp}{sp}\binom{2mp^r-2sp}{mp^r-sp}}{\binom{2s}{s}\binom{2mp^{r-1}-2s}{mp^{r-1}-s}} = \frac{\prod\limits_{\substack{i=1\\ p\nmid i}}^{mp^r} \frac{2mp^r-i}{i} \prod\limits_{\substack{i=1\\ p\nmid i}}^{sp} (\frac{mp^r-i}{i})^2}{\prod\limits_{\substack{i=1\\ p\nmid i}}^{2sp} \frac{2mp^r-i}{i}} \equiv \frac{\Big(-2mp^r \sum\limits_{\substack{i=1\\ p\nmid i}}^{mp^r} \frac{1}{i} + 1\Big)\Big(-2mp^r \sum\limits_{\substack{i=1\\ p\nmid i}}^{sp} \frac{1}{i} + 1\Big)}{-2mp^r \sum\limits_{\substack{i=1\\ p\nmid i}}^{2sp} \frac{1}{i} + 1}$$
$$\equiv \frac{-2mp^r \sum\limits_{\substack{i=1\\ p\nmid i}}^{mp^r} \frac{1}{i} - 2mp^r \sum\limits_{\substack{i=1\\ p\nmid i}}^{sp} \frac{1}{i} + 1}{-2mp^r \sum\limits_{\substack{i=1\\ p\nmid i}}^{2sp} \frac{1}{i} + 1} \equiv 1 \pmod{p^{r+\min\{l+1,r\}}}.$$

Thus,

$$\binom{2sp}{sp}\binom{2mp^r-2sp}{mp^r-sp} \equiv \binom{2s}{s}\binom{2mp^{r-1}-2s}{mp^{r-1}-s} \pmod{p^{r+\min\{l+1,r\}}}.$$

If $l \geq r-1$, then $r+l+1 \geq 2r$ and so

$$\binom{mp^r}{sp}\binom{2sp}{sp}\binom{2mp^r-2sp}{mp^r-sp} \equiv \binom{mp^{r-1}}{s}\binom{2s}{s}\binom{2mp^{r-1}-2s}{mp^{r-1}-s} \pmod{p^{2r}}.$$

If $1 \leq l < r-1$, then $\binom{mp^{r-1}}{s} \equiv 0 \pmod{p^{r-1-l}}$ and so

$$\binom{mp^r}{sp}\binom{2sp}{sp}\binom{2mp^r-2sp}{mp^r-sp} \equiv \binom{mp^{r-1}}{s}\binom{2s}{s}\binom{2mp^{r-1}-2s}{mp^{r-1}-s} \pmod{p^{2r}}.$$

Now the proof is complete.

Lemma 2.15 ([B]). *Let p be an odd prime. Suppose $n = n_1 p + n_0$ and $k = k_1 p + k_0$ with $k_1, n_1 \in \mathbb{Z}^+$ and $k_0, n_0 \in \{0, 1, \ldots, p-1\}$. Then*

$$\binom{n}{k} \equiv \binom{n_1}{k_1}\Big[(1+n_1)\binom{n_0}{k_0} - (n_1+k_1)\binom{n_0-p}{k_0} - k_1\binom{n_0-p}{k_0+p}\Big] \pmod{p^2}.$$

Lemma 2.16. *Let p be an odd prime. Then*

$$\sum_{t=0}^{(p-1)/2}(-1)^t\Big[\binom{\frac{p-1}{2}+t}{t}-\binom{p+\frac{p-1}{2}+t}{p+t}\Big]\binom{-\frac{1}{2}}{t}^2\equiv 0 \pmod{p^2}.$$

Proof. From Lemma 2.2 we have

$$\begin{aligned}\binom{\frac{p-1}{2}-p}{p+t}&=\binom{-\frac{p+1}{2}}{p+t}=(-1)^{t+1}\binom{p+\frac{p+1}{2}+t-1}{p+t}\equiv(-1)^{t+1}\binom{\frac{p+1}{2}+t-1}{t}\\&\equiv(-1)^{t+1}\binom{\frac{1}{2}+t-1}{t}=-\binom{-\frac{1}{2}}{t}\equiv-\binom{\frac{p-1}{2}-p}{t}\pmod p\end{aligned}$$

and so $\binom{\frac{p-1}{2}-p}{t}+\binom{\frac{p-1}{2}-p}{p+t}=(-1)^t\Big(\binom{\frac{p-1}{2}+t}{t}-\binom{p+\frac{p-1}{2}+t}{p+t}\Big)\equiv 0\pmod p$.

We first assume $p\equiv 1\pmod 4$. Applying Lemma 2.15 we get

$$\begin{aligned}\binom{\frac{3(p-1)}{4}}{\frac{p-1}{4}}-\binom{p+\frac{3(p-1)}{4}}{p+\frac{p-1}{4}}&=\binom{\frac{3(p-1)}{4}}{\frac{p-1}{2}}-\binom{p+\frac{3(p-1)}{4}}{\frac{p-1}{2}}\\&\equiv\binom{\frac{3(p-1)}{4}}{\frac{p-1}{2}}-\Big(2\binom{\frac{3(p-1)}{4}}{\frac{p-1}{2}}-\binom{\frac{3(p-1)}{4}-p}{\frac{p-1}{2}}\Big)\\&=-\binom{\frac{3(p-1)}{4}}{\frac{p-1}{2}}+(-1)^{\frac{p-1}{2}}\binom{\frac{3(p-1)}{4}}{\frac{p-1}{2}}=0\pmod{p^2},\end{aligned}$$

and

$$\begin{aligned}&\binom{\frac{p-1}{2}+t}{t}-\binom{p+\frac{p-1}{2}+t}{p+t}+\binom{p-1-t}{\frac{p-1}{2}-t}-\binom{p+p-1-t}{p+\frac{p-1}{2}-t}\\&=\binom{\frac{p-1}{2}+t}{\frac{p-1}{2}}-\binom{p+\frac{p-1}{2}+t}{\frac{p-1}{2}}+\binom{p-1-t}{\frac{p-1}{2}}-\binom{p+p-1-t}{\frac{p-1}{2}}\\&\equiv\binom{\frac{p-1}{2}+t}{\frac{p-1}{2}}-\Big(2\binom{\frac{p-1}{2}+t}{\frac{p-1}{2}}-\binom{\frac{p-1}{2}-p+t}{\frac{p-1}{2}}\Big)+\binom{p-1-t}{\frac{p-1}{2}}-\Big(2\binom{p-1-t}{\frac{p-1}{2}}-\binom{-1-t}{\frac{p-1}{2}}\Big)\\&=-\binom{\frac{p-1}{2}+t}{\frac{p-1}{2}}+(-1)^{\frac{p-1}{2}}\binom{p-1-t}{\frac{p-1}{2}}-\binom{p-1-t}{\frac{p-1}{2}}+(-1)^{\frac{p-1}{2}}\binom{\frac{p-1}{2}+t}{\frac{p-1}{2}}\\&=0\pmod{p^2}.\end{aligned}$$

Also,

$$\begin{aligned}(-1)^t\binom{-\frac{1}{2}}{t}^2-(-1)^{\frac{p-1}{2}-t}\binom{-\frac{1}{2}}{\frac{p-1}{2}-t}^2&\equiv(-1)^t\binom{-\frac{1}{2}}{t}^2-(-1)^t\binom{\frac{p-1}{2}}{\frac{p-1}{2}-t}^2\\&\equiv(-1)^t\binom{-\frac{1}{2}}{t}^2-(-1)^t\binom{-\frac{1}{2}}{t}^2=0\pmod p.\end{aligned}$$

Hence

$$\sum_{t=0}^{(p-1)/2}(-1)^t\Big[\binom{\frac{p-1}{2}+t}{t}-\binom{p+\frac{p-1}{2}+t}{p+t}\Big]\binom{-\frac{1}{2}}{t}^2$$

$$
\begin{aligned}
&= \sum_{t=0}^{(p-5)/4} (-1)^t \binom{-\frac12}{t}^2 \Big[\binom{\frac{p-1}{2}+t}{t} - \binom{p+\frac{p-1}{2}+t}{p+t}\Big] \\
&\quad + (-1)^{\frac{p-1}{4}} \binom{-\frac12}{\frac{p-1}{4}}^2 \Big[\binom{\frac{3(p-1)}{4}}{\frac{p-1}{4}} - \binom{p+\frac{3(p-1)}{4}}{p+\frac{p-1}{4}}\Big] \\
&\quad + \sum_{t=0}^{(p-5)/4} (-1)^{\frac{p-1}{2}-t} \binom{-\frac12}{\frac{p-1}{2}-t}^2 \Big[\binom{p-1-t}{\frac{p-1}{2}-t} - \binom{p+p-1-t}{p+\frac{p-1}{2}-t}\Big] \\
&\equiv \sum_{t=0}^{(p-5)/4} (-1)^t \binom{-\frac12}{t}^2 \Big[\binom{\frac{p-1}{2}+t}{t} - \binom{p+\frac{p-1}{2}+t}{p+t}\Big] \\
&\quad + (-1)^{\frac{p-1}{4}} \binom{\frac{3(p-1)}{4}}{\frac{p-1}{4}} \Big[\binom{\frac{3(p-1)}{4}}{\frac{p-1}{4}} - \binom{p+\frac{3(p-1)}{4}}{p+\frac{p-1}{4}}\Big] \\
&\quad - \sum_{t=0}^{(p-5)/4} (-1)^{\frac{p-1}{2}-t} \binom{-\frac12}{\frac{p-1}{2}-t}^2 \Big[\binom{\frac{p-1}{2}+t}{t} - \binom{p+\frac{p-1}{2}+t}{p+t}\Big] \\
&\equiv \sum_{t=0}^{(p-5)/4} \Big[(-1)^t \binom{-\frac12}{t}^2 - (-1)^{\frac{p-1}{2}-t} \binom{-\frac12}{\frac{p-1}{2}-t}^2\Big] \Big[\binom{\frac{p-1}{2}+t}{t} - \binom{p+\frac{p-1}{2}+t}{p+t}\Big] \\
&\quad + (-1)^{\frac{p-1}{4}} \binom{\frac{3(p-1)}{4}}{\frac{p-1}{4}} \Big[\binom{\frac{3(p-1)}{4}}{\frac{p-1}{4}} - \binom{p+\frac{3(p-1)}{4}}{p+\frac{p-1}{4}}\Big] \equiv 0 \pmod{p^2}.
\end{aligned}
$$

Thus the result is true for $p \equiv 1 \pmod 4$.

Now we assume $p \equiv 3 \pmod 4$. By Lemma 2.15,

$$\binom{\frac{p-1}{2}+t}{t} - \binom{p+\frac{p-1}{2}+t}{p+t} \equiv -\Big(\binom{\frac{p-1}{2}+t}{\frac{p-1}{2}} + \binom{p-1-t}{\frac{p-1}{2}}\Big) \pmod{p^2}.$$

As

$$\binom{\frac{p-1}{2}+t}{\frac{p-1}{2}} + \binom{p-1-t}{\frac{p-1}{2}} \equiv \binom{\frac{p-1}{2}+t}{\frac{p-1}{2}} + \binom{-1-t}{\frac{p-1}{2}} = \binom{\frac{p-1}{2}+t}{\frac{p-1}{2}} + (-1)^{\frac{p-1}{2}} \binom{t+\frac{p-1}{2}}{\frac{p-1}{2}} = 0 \pmod p$$

and

$$
\begin{aligned}
&(-1)^t \binom{-\frac12}{t}^2 + (-1)^{\frac{p-1}{2}-t} \binom{-\frac12}{\frac{p-1}{2}-t}^2 \equiv (-1)^t \binom{-\frac12}{t}^2 - (-1)^t \binom{\frac{p-1}{2}}{\frac{p-1}{2}-t}^2 \\
&\equiv (-1)^t \binom{-\frac12}{t}^2 - (-1)^t \binom{-\frac12}{t}^2 = 0 \pmod p,
\end{aligned}
$$

we obtain

$$
\begin{aligned}
&\sum_{t=0}^{(p-1)/2} (-1)^t \Big[\binom{\frac{p-1}{2}+t}{t} - \binom{p+\frac{p-1}{2}+t}{p+t}\Big] \binom{-\frac12}{t}^2 \\
&\equiv - \sum_{t=0}^{(p-3)/4} \Big[\binom{\frac{p-1}{2}+t}{\frac{p-1}{2}} + \binom{p-1-t}{\frac{p-1}{2}}\Big] \Big[(-1)^t \binom{-\frac12}{t}^2 + (-1)^{\frac{p-1}{2}-t} \binom{-\frac12}{\frac{p-1}{2}-t}^2\Big] \\
&\equiv 0 \pmod{p^2}.
\end{aligned}
$$

Hence the result is also true in this case. The proof is now complete.

3. Congruences for $S_{mp^r} \pmod{p^{r+2}}$, $S_{mp^r-1} \pmod{p^r}$ and $S_{mp^r+1} \pmod{p^{2r}}$

Theorem 3.1. *Let p be an odd prime and $n \in \mathbb{Z}^+$. Then*

$$S_{np} - S_n \equiv \begin{cases} 8n^2 S_{n-1}(-1)^{\frac{p-1}{2}} p^2 E_{p-3} \pmod{p^3} & \text{if } p > 3 \text{ and } p \nmid n, \\ 9(n-1)S_n \pmod{p^3} & \text{if } p = 3 \text{ and } 3 \nmid n, \\ 0 \pmod{p^{3+\mathrm{ord}_p n}} & \text{if } p \mid n. \end{cases}$$

Proof. Set $r = \mathrm{ord}_p(np)$. Then

$$\begin{aligned} S_{np} &= \sum_{k=0}^{np} \binom{np}{k}\binom{2k}{k}\binom{2(np-k)}{np-k} \\ &= \sum_{s=0}^{n} \binom{np}{sp}\binom{2sp}{sp}\binom{2(n-s)p}{(n-s)p} + \sum_{t=1}^{p-1}\sum_{s=0}^{n-1} \binom{np}{sp+t}\binom{2(sp+t)}{sp+t}\binom{2(np-sp-t)}{np-sp-t}. \end{aligned}$$

If $p > 3$ or if $p = 3$ and $3 \mid n$, using Lemmas 2.10 and 2.11 we see that $\binom{np}{sp} \equiv \binom{n}{s} \pmod{p^{r+2}}$ and

$$\binom{n}{s}\binom{2sp}{sp}\binom{2(n-s)p}{(n-s)p} \equiv \binom{n}{s}\binom{2s}{s}\binom{2(n-s)}{n-s} \pmod{p^{r+2}}.$$

Thus,

$$\begin{aligned} &\sum_{s=0}^{n} \binom{np}{sp}\binom{2sp}{sp}\binom{2(n-s)p}{(n-s)p} \\ &\equiv \sum_{s=0}^{n} \binom{n}{s}\binom{2sp}{sp}\binom{2(n-s)p}{(n-s)p} \equiv \sum_{s=0}^{n} \binom{n}{s}\binom{2s}{s}\binom{2(n-s)}{n-s} = S_n \pmod{p^{r+2}}. \end{aligned}$$

Hence

$$\begin{aligned} &S_{np} - S_n \\ &\equiv \sum_{t=1}^{p-1}\sum_{s=0}^{n-1} \frac{np}{sp+t}\binom{(n-1)p+p-1}{sp+t-1}\binom{2sp+2t}{sp+t}\binom{2(n-1-s)p+2(p-t)}{(n-1-s)p+p-t} \pmod{p^{r+2}}. \end{aligned}$$

For $t \in \{1, 2, \ldots, \frac{p-1}{2}\}$ we have $\frac{p}{2} < p - t < p$. By Lemma 2.6,

$$\binom{2(n-1-s)p+2(p-t)}{(n-1-s)p+p-t} \equiv (2(n-1-s)+1)\binom{2(n-1-s)}{n-1-s}\binom{2(p-t)}{p-t} \pmod{p^2}.$$

By Lemma 2.2, $\binom{2sp+2t}{sp+t} \equiv \binom{2s}{s}\binom{2t}{t} \pmod p$. Thus, applying Lemma 2.4 we see that

$$\begin{aligned} &\binom{2sp+2t}{sp+t}\binom{2(n-1-s)p+2(p-t)}{(n-1-s)p+p-t} \\ &\equiv \binom{2s}{s}\binom{2t}{t}(2(n-1-s)+1)\binom{2(n-1-s)}{n-1-s}\binom{2(p-t)}{p-t} \end{aligned}$$

$$\equiv -(2(n-1-s)+1)\binom{2s}{s}\binom{2(n-1-s)}{n-1-s}\frac{2p}{t} \pmod{p^2}.$$

For $t \in \{\frac{p+1}{2}, \dots, p-1\}$ we have $1 \le p-t < \frac{p}{2}$. By Lemma 2.6,

$$\binom{2sp+2t}{sp+t} \equiv (2s+1)\binom{2s}{s}\binom{2t}{t} \pmod{p^2}.$$

By Lemma 2.2,

$$\binom{2(n-1-s)p+2(p-t)}{(n-1-s)p+p-t} \equiv \binom{2(n-1-s)}{n-1-s}\binom{2(p-t)}{p-t} \pmod{p}.$$

Thus, applying Lemma 2.4 we get

$$\begin{aligned}
&\binom{2sp+2t}{sp+t}\binom{2(n-1-s)p+2(p-t)}{(n-1-s)p+p-t}\\
&\equiv (2s+1)\binom{2s}{s}\binom{2t}{t}\binom{2(n-1-s)}{n-1-s}\binom{2(p-t)}{p-t}\\
&\equiv (2s+1)\binom{2s}{s}\binom{2(n-1-s)}{n-1-s}\frac{2p}{t} \pmod{p^2}.
\end{aligned}$$

Hence

$$\begin{aligned}
&S_{np}-S_n\\
&\equiv \sum_{t=1}^{(p-1)/2}\sum_{s=0}^{n-1}\frac{np}{sp+t}\binom{(n-1)p+p-1}{sp+t-1}\binom{2sp+2t}{sp+t}\binom{2(n-1-s)p+2(p-t)}{(n-1-s)p+p-t}\\
&\quad+\sum_{t=(p+1)/2}^{p-1}\sum_{s=0}^{n-1}\frac{np}{sp+t}\binom{(n-1)p+p-1}{sp+t-1}\binom{2sp+2t}{sp+t}\binom{2(n-1-s)p+2(p-t)}{(n-1-s)p+p-t}\\
&\equiv -\sum_{t=1}^{(p-1)/2}\sum_{s=0}^{n-1}\frac{np}{sp+t}\binom{(n-1)p+p-1}{sp+t-1}(2(n-1-s)+1)\binom{2s}{s}\binom{2(n-1-s)}{n-1-s}\frac{2p}{t}\\
&\quad+\sum_{t=(p+1)/2}^{p-1}\sum_{s=0}^{n-1}\frac{np}{sp+t}\binom{(n-1)p+p-1}{sp+t-1}(2s+1)\binom{2s}{s}\binom{2(n-1-s)}{n-1-s}\frac{2p}{t}\\
&\equiv -\sum_{t=1}^{(p-1)/2}\sum_{s=0}^{n-1}\frac{np}{t}\binom{n-1}{s}\binom{p-1}{t-1}(2(n-1-s)+1)\binom{2s}{s}\binom{2(n-1-s)}{n-1-s}\frac{2p}{t}\\
&\quad+\sum_{t=(p+1)/2}^{p-1}\sum_{s=0}^{n-1}\frac{np}{t}\binom{n-1}{s}\binom{p-1}{t-1}(2s+1)\binom{2s}{s}\binom{2(n-1-s)}{n-1-s}\frac{2p}{t}\\
&\equiv 2np^2\sum_{s=0}^{n-1}(2(n-1-s)+1)\binom{n-1}{s}\binom{2s}{s}\binom{2(n-1-s)}{n-1-s}\sum_{t=1}^{(p-1)/2}\frac{(-1)^t}{t^2}\\
&\quad-2np^2\sum_{s=0}^{n-1}(2s+1)\binom{n-1}{s}\binom{2s}{s}\binom{2(n-1-s)}{n-1-s}\sum_{t=(p+1)/2}^{p-1}\frac{(-1)^t}{t^2}
\end{aligned}$$

$$\equiv 2np^2\sum_{s=0}^{n-1}(2s+1)\binom{n-1}{s}\binom{2s}{s}\binom{2(n-1-s)}{n-1-s}$$
$$\times\Big(\sum_{t=1}^{(p-1)/2}\frac{(-1)^t}{t^2}-\sum_{t=(p+1)/2}^{p-1}\frac{(-1)^t}{t^2}\Big)\pmod{p^{r+2}}.$$

By Lemma 2.7,

$$\begin{aligned}&\sum_{s=0}^{n-1}(2s+1)\binom{n-1}{s}\binom{2s}{s}\binom{2(n-1-s)}{n-1-s}\\&=S_{n-1}+2\sum_{s=1}^{n-1}s\binom{n-1}{s}\binom{2s}{s}\binom{2(n-1-s)}{n-1-s}\\&=S_{n-1}+2(n-1)\sum_{s=1}^{n-1}\binom{n-2}{s-1}\binom{2s}{s}\binom{2(n-1-s)}{n-1-s}\\&=S_{n-1}+(n-1)S_{n-1}=nS_{n-1}.\end{aligned}$$

Note that $B_{p-2}=0$ and $E_{2n}=2^{2n}E_{2n}(\frac{1}{2})$. From Lemma 2.5 we see that

$$\sum_{k=1}^{(p-1)/2}\frac{(-1)^k}{k^2}\equiv\frac{1}{2}(-1)^{\frac{p-1}{2}}E_{p-3}\Big(\frac{1}{2}\Big)\equiv 2(-1)^{\frac{p-1}{2}}E_{p-3}\pmod p.$$

Thus,

$$\begin{aligned}&\sum_{k=1}^{(p-1)/2}\frac{(-1)^k}{k^2}-\sum_{k=(p+1)/2}^{p-1}\frac{(-1)^k}{k^2}\\&=\sum_{k=1}^{(p-1)/2}\frac{(-1)^k}{k^2}-\sum_{k=1}^{(p-1)/2}\frac{(-1)^{p-k}}{(p-k)^2}\equiv 2\sum_{k=1}^{(p-1)/2}\frac{(-1)^k}{k^2}\equiv 4(-1)^{\frac{p-1}{2}}E_{p-3}\pmod p.\end{aligned}$$

Now from the above we deduce that $S_{np}-S_n\equiv 2np^2\cdot nS_{n-1}\cdot 4(-1)^{\frac{p-1}{2}}E_{p-3}\pmod{p^{r+2}}$. This yields the result in this case.

Now assume $3\nmid n$. By Lemmas 2.9 and 2.11,

$$\binom{3n}{3s}\equiv\binom{n}{s}(1+9ns^2-9s)\pmod{27},$$

and

$$\begin{aligned}\binom{3n}{3s}\binom{6s}{3s}\binom{6(n-s)}{3(n-s)}&\equiv\binom{n}{s}\binom{2s}{s}\binom{2(n-s)}{n-s}(1+9n)(1+9ns^2-9s)\\&\equiv\binom{n}{s}\binom{2s}{s}\binom{2(n-s)}{n-s}(1+9n+9ns^2-9s)\pmod{27}.\end{aligned}$$

Thus,

$$\sum_{s=0}^{n}\binom{3n}{3s}\binom{6s}{3s}\binom{6(n-s)}{3(n-s)}\equiv(1+9n)S_n+9\sum_{s=0}^{n}(ns^2-s)\binom{n}{s}\binom{2s}{s}\binom{2(n-s)}{n-s}\pmod{27}$$

and so

$$\begin{aligned}
&S_{3n}-S_n\\
&\equiv \sum_{t=1}^{2}\sum_{s=0}^{n-1}\frac{3n}{3s+t}\binom{3(n-1)+3-1}{3s+t-1}\binom{6s+2t}{3s+t}\binom{6(n-1-s)+2(3-t)}{3(n-1-s)+3-t}\\
&+9nS_n+9\sum_{s=0}^{n}(ns^2-s)\binom{n}{s}\binom{2s}{s}\binom{2(n-s)}{n-s}\\
&\equiv 2n3^2nS_{n-1}(-5/4)+9nS_n+9\sum_{s=0}^{n}(ns^2-s)\binom{n}{s}\binom{2s}{s}\binom{2(n-s)}{n-s}\\
&\equiv 9nS_n-9S_{n-1}+9\sum_{s=0}^{n}(ns^2-s)\binom{n}{s}\binom{2s}{s}\binom{2(n-s)}{n-s} \pmod{27}.
\end{aligned}$$

By (1.3), for $n\equiv 2 \pmod 3$ we have

$$S_n+S_{n-1}\equiv 4(3n(n+1)+1)S_n-32n^2S_{n-1}=(n+1)^2S_{n+1}\equiv 0 \pmod 3,$$

for $n\equiv 1 \pmod 3$ we have

$$S_n-S_{n-1}\equiv n^2S_n-4(3n(n-1)+1)S_{n-1}=-32(n-1)^2S_{n-2}\equiv 0 \pmod 3.$$

Thus,

$$S_n\equiv\Big(\frac{n}{3}\Big)S_{n-1} \pmod 3 \quad\text{for}\quad n\not\equiv 0 \pmod 3. \tag{3.1}$$

Applying (3.1) and (2.2) we have

$$\begin{aligned}
S_{3n}-S_n&\equiv 9(nS_n-S_{n-1})+9\sum_{s=0}^{n}(ns^2-s)\binom{n}{s}\binom{2s}{s}\binom{2(n-s)}{n-s}\\
&\equiv 9\sum_{s=0}^{n}(ns^2-s)\binom{n}{s}\binom{2s}{s}\binom{2(n-s)}{n-s}\\
&=9\sum_{s=0}^{n}\Big(ns(s-1)+\frac{n-1}{2}(2s-n)+\frac{n(n-1)}{2}\Big)\binom{n}{s}\binom{2s}{s}\binom{2(n-s)}{n-s}\\
&=9n\sum_{s=0}^{n}s(s-1)\binom{n}{s}\binom{2s}{s}\binom{2(n-s)}{n-s}+\frac{9n(n-1)}{2}S_n \pmod{27}.
\end{aligned}$$

If $s=3k+2$ for some nonnegative integer k, using Lemma 2.2 we find that $\binom{2s}{s}=\binom{3(2k+1)+1}{3k+2}\equiv\binom{2k+1}{k}\binom{1}{2}=0 \pmod 3$. Thus, $3\mid s(s-1)\binom{2s}{s}$ for any nonnegative integer s. Hence, from the above we deduce that

$$S_{3n}-S_n\equiv\frac{9n(n-1)}{2}S_n\equiv 9(n-n^2)S_n\equiv 9(n-1)S_n \pmod{27}.$$

This completes the proof.

Corollary 3.1. *Let $p>3$ be a prime. Then*

$$S_p\equiv 4+8(-1)^{\frac{p-1}{2}}p^2E_{p-3} \pmod{p^3},$$

$$S_{2p} \equiv 20 + 128(-1)^{\frac{p-1}{2}} p^2 E_{p-3} \pmod{p^3},$$
$$S_{3p} \equiv 112 + 1440(-1)^{\frac{p-1}{2}} p^2 E_{p-3} \pmod{p^3}.$$

Remark 3.1. *Let p be an odd prime and $m, r \in \mathbb{Z}^+$. Then $S_{mp^r} \equiv S_{mp^{r-1}} \pmod{p^{2r}}$.*

Proof. $S_{mp^r} = \sum_{s=0}^{mp^{r-1}} \binom{mp^r}{sp}\binom{2sp}{sp}\binom{2mp^r-2sp}{mp^r-sp} + \sum_{s=0}^{mp^{r-1}-1}\sum_{t=1}^{p-1} \binom{mp^r}{sp+t}\binom{2sp+2t}{sp+t}\binom{2mp^r-2sp-2t}{mp^r-sp-t}$. Applying Lemmas 2.12 and 2.14 we obtain

$$S_{mp^r} \equiv \sum_{s=0}^{mp^{r-1}} \binom{mp^{r-1}}{s}\binom{2s}{s}\binom{2mp^{r-1}-2s}{mp^{r-1}-s} = S_{mp^{r-1}} \pmod{p^{2r}}.$$

Lemma 3.1. *Let $m, n \in \mathbb{Z}^+$. Then*

$$S_{mn+1} \equiv 4(mn+1)S_{mn} \pmod{m^2n^2}.$$

Proof. By (1.3),

$$(mn+1)^2 S_{mn+1} = 4(3mn(mn+1)+1)S_{mn} - 32m^2n^2 S_{mn-1}.$$

Thus,

$$(1+2mn)S_{mn+1} \equiv (mn+1)^2 S_{mn+1} \equiv 4(3mn+1)S_{mn} \pmod{m^2n^2}$$

and so

$$S_{mn+1} \equiv \frac{4(1+3mn)}{1+2mn} S_{mn} \equiv 4(1+3mn)(1-2mn)S_{mn} \equiv 4(1+mn)S_{mn} \pmod{m^2n^2}$$

as asserted.

Theorem 3.2. *Let p be an odd prime, and $m, r \in \mathbb{Z}^+$. Then*

$$S_{mp^r+1} \equiv 4(mp^r+1)S_{mp^{r-1}} \pmod{p^{2r}}.$$

Proof. As $\mathrm{ord}_p(m^2p^{2r}) \geq 2r$, from Lemma 3.1 and Remark 3.1 we see that

$$S_{mp^r+1} \equiv 4(mp^r+1)S_{mp^r} \equiv 4(mp^r+1)S_{mp^{r-1}} \pmod{p^{2r}}.$$

This completes the proof.

Theorem 3.3. *Let p be an odd prime and $m, r \in \mathbb{Z}^+$. Then*

$$S_{mp^r-1} \equiv (-1)^{\frac{p-1}{2}} S_{mp^{r-1}-1} \pmod{p^r}.$$

Proof. It is clear that

$$\begin{aligned}
S_{mp^r-1} &= \sum_{s=0}^{mp^{r-1}-1}\sum_{t=0}^{p-1} \binom{2sp+2t}{sp+t}\binom{mp^r-1}{sp+t}\binom{2(mp^r-1-sp-t)}{mp^r-1-sp-t} \\
&= \sum_{s=0}^{mp^{r-1}-1}\sum_{\substack{t=0\\ t\neq(p-1)/2}}^{p-1} \binom{2sp+2t}{sp+t}\binom{mp^r-1}{sp+t}\binom{2(mp^r-1-sp-t)}{mp^r-1-sp-t} \\
&\quad + \sum_{s=0}^{mp^{r-1}-1} \binom{2sp+p-1}{sp+\frac{p-1}{2}}\binom{mp^r-1}{sp+\frac{p-1}{2}}\binom{2(mp^r-1-sp-\frac{p-1}{2})}{mp^r-1-sp-\frac{p-1}{2}}.
\end{aligned}$$

Using Lemma 2.8 we see that

$$\binom{mp^r-1}{sp+t} \equiv \binom{mp^{r-1}-1}{s}(-1)^t \pmod{p^r}.$$

For $t \ne \frac{p-1}{2}$ applying Lemma 2.12 we obtain

$$\begin{aligned}&\binom{2sp+2t}{sp+t}\binom{2(mp^r-1-sp-t)}{mp^r-1-sp-t}\\ &= \binom{2sp+2t}{sp+t}\binom{2(mp^r-sp-t)}{mp^r-sp-t}\frac{(mp^r-sp-t)^2}{(2mp^r-1-2sp-2t)2(mp^r-sp-t)}\\ &\equiv \binom{2sp+2t}{sp+t}\binom{2(mp^r-sp-t)}{mp^r-sp-t}\frac{sp+t}{2(2sp+2t+1)} \equiv 0 \pmod{p^r}.\end{aligned}$$

For $t = \frac{p-1}{2}$ using Lemma 2.13 we deduce that

$$\begin{aligned}S_{mp^r-1} &\equiv (-1)^{\frac{p-1}{2}} \sum_{s=0}^{mp^{r-1}-1}\binom{2s}{s}\binom{mp^{r-1}-1}{s}\binom{2(mp^{r-1}-1-s)}{mp^{r-1}-1-s}\\ &= (-1)^{\frac{p-1}{2}} S_{mp^{r-1}-1} \pmod{p^r}.\end{aligned}$$

So the theorem is proved.

Theorem 3.4. *Let p be an odd prime and $n \in \mathbb{Z}^+$. Then*

$$S_{np+1} \equiv \begin{cases} (4+12n-9n^2)S_n \pmod{p^3} & \text{if } p=3,\\ 4(np+1)S_n + 32n^2S_{n-1}(-1)^{\frac{p-1}{2}}(E_{p-3}-1)p^2 \pmod{p^3} & \text{if } p>3.\end{cases}$$

Proof. By (1.3), $(np+1)^2S_{np+1} = 4(3np(np+1)+1)S_{np} - 32n^2p^2S_{np-1}$. Thus, applying Theorems 3.1 and 3.3 we see that for $p > 3$,

$$\begin{aligned}(np+1)^2S_{np+1} &\equiv 4(3n^2p^2+3np+1)(S_n + 8n^2S_{n-1}(-1)^{\frac{p-1}{2}}p^2E_{p-3}) - 32n^2p^2(-1)^{\frac{p-1}{2}}S_{n-1}\\ &\equiv 4(3n^2p^2+3np+1)S_n + 32n^2S_{n-1}(-1)^{\frac{p-1}{2}}(E_{p-3}-1)p^2 \pmod{p^3},\end{aligned}$$

and for $p = 3$,

$$\begin{aligned}(3n+1)^2S_{3n+1} &= 4(9n(3n+1)+1)S_{3n} - 32n^2\cdot 9S_{3n-1}\\ &\equiv 4(9n+1)(1-9n(n-1))S_n - 9n^2S_{n-1}\\ &\equiv 4(1-9n(n+1))S_n - 9n^2S_{n-1} \pmod{27}.\end{aligned} \tag{3.2}$$

Since

$$\frac{1}{(np+1)^2} = \frac{(n^2p^2-np+1)^2}{\big((np)^3+1\big)^2} \equiv (n^2p^2-np+1)^2 \equiv 3n^2p^2-2np+1 \pmod{p^3}, \tag{3.3}$$

from the above we deduce that for $p > 3$,

$$\begin{aligned}S_{np+1} &\equiv \frac{4(3n^2p^2+3np+1)S_n + 32n^2S_{n-1}(-1)^{\frac{p-1}{2}}(E_{p-3}-1)p^2}{(np+1)^2}\\ &\equiv 4(S_n + 3npS_n + n^2p^2(3S_n + 8S_{n-1}(-1)^{\frac{p-1}{2}}(E_{p-3}-1))(3n^2p^2-2np+1)\\ &\equiv 4(np+1)S_n + 32n^2S_{n-1}(-1)^{\frac{p-1}{2}}(E_{p-3}-1)p^2 \pmod{p^3}.\end{aligned}$$

Now assume $p=3$. If $3\mid n$, from (3.2) and (3.3) we deduce that

$$S_{3n+1}\equiv \frac{4S_n}{(3n+1)^2}\equiv 4(1-6n)S_n\equiv (4+12n-9n^2)S_n \pmod{27}.$$

If $3\nmid n$, then $S_{n-1}\equiv (\frac{n}{3})S_n \pmod 3$ by (3.1). Hence, from (3.2) and (3.3) we deduce that

$$\begin{aligned}S_{3n+1}&\equiv \frac{4S_n-9n((n+1)S_n+nS_{n-1})}{(3n+1)^2}\equiv \frac{4S_n-9(n+1+(\frac{n}{3}))S_n}{(3n+1)^2}\\&\equiv (4-9(2n+1))S_n(1-6n)\equiv (12n-5)S_n\equiv (4+12n-9n^2)S_n \pmod{27}.\end{aligned}$$

Summarizing the above proves the theorem.

Corollary 3.2. *Let $p>3$ be a prime. Then*

$$S_{p+1}\equiv 16+16p+32(-1)^{\frac{p-1}{2}}(E_{p-3}-1)p^2 \pmod{p^3}.$$

Proof. Taking $n=1$ in Theorem 3.4 we obtain the result.

Theorem 3.5. *Let p be a prime with $p\equiv 5,7 \pmod 8$. Then*

$$S_{\frac{p^2-1}{2}}\equiv 0 \pmod{p^2} \quad \text{and} \quad f_{\frac{p^2-1}{2}}\equiv 0 \pmod{p^2}.$$

Proof. For $\frac{p-1}{2}<t<p$, from Lemma 2.2 we see that $p\mid\binom{2sp+2t}{sp+t}$ and $p\mid\binom{\frac{p-1}{2}p+\frac{p-1}{2}}{sp+t}$. So

$$f_{\frac{p^2-1}{2}}\equiv \sum_{s=0}^{\frac{p-1}{2}}\sum_{t=0}^{\frac{p-1}{2}}\binom{\frac{p-1}{2}p+\frac{p-1}{2}}{sp+t}^3 \pmod{p^2}$$

and

$$S_{\frac{p^2-1}{2}}\equiv \sum_{s=0}^{\frac{p-1}{2}}\sum_{t=0}^{\frac{p-1}{2}}\binom{\frac{p-1}{2}p+\frac{p-1}{2}}{sp+t}\binom{2sp+2t}{sp+t}\binom{(p-1-2s)p+p-1-2t}{(\frac{p-1}{2}-s)p+\frac{p-1}{2}-t} \pmod{p^2}.$$

For $k\in\mathbb{Z}^+$ set $H_k=1+\frac12+\cdots+\frac1k$. For $k\in\{1,\ldots,p-1\}$ we see that

$$\begin{aligned}\binom{p-1}{k}&=\frac{(p-1)(p-2)\cdots(p-k)}{k!}\equiv\frac{(-1)(-2)\cdots(-k)\big(1+\sum_{i=1}^{k}\frac{p}{-i}\big)}{k!}\\&=(-1)^k(1-pH_k) \pmod{p^2}\end{aligned}$$

and so $\frac{1}{\binom{p-1}{2s}}\equiv 1+pH_{2s} \pmod{p^2}$ for $s=1,2,\ldots,\frac{p-1}{2}$. It is well known that $H_{\frac{p-1}{2}}\equiv -\frac{2^p-2}{p} \pmod p$. Thus,

$$\binom{p-1}{\frac{p-1}{2}}\equiv (-1)^{\frac{p-1}{2}}(1-pH_{\frac{p-1}{2}})\equiv (-1)^{\frac{p-1}{2}}(2^p-1) \pmod{p^2}.$$

Applying (2.4) and Lemma 2.8 we have

$$\begin{aligned}
&\binom{2sp+2t}{sp+t}\binom{p^2-1-2sp-2t}{\frac{p^2-1}{2}-sp-t}\\
&= \frac{\binom{p^2-1}{\frac{p^2-1}{2}}\binom{\frac{p-1}{2}p+\frac{p-1}{2}}{sp+t}^2}{\binom{p^2-1}{2sp+2t}} = \frac{\binom{p-1}{\frac{p-1}{2}}\binom{\frac{p-1}{2}p+\frac{p-1}{2}}{sp+t}^2}{\binom{p-1}{2s}} \frac{\prod\limits_{\substack{i=1\\ p\nmid i}}^{(p^2-1)/2} \frac{p^2-i}{i}}{\prod\limits_{\substack{i=1\\ p\nmid i}}^{2sp+2t} \frac{p^2-i}{i}}\\
&\equiv (-1)^{\frac{p-1}{2}}(2^p-1)(1+pH_{2s})\binom{\frac{p-1}{2}p+\frac{p-1}{2}}{sp+t}^2\\
&\equiv \Big((-1)^{\frac{p-1}{2}}(2^p-1)+(-1)^{\frac{p-1}{2}}pH_{2s}\Big)\binom{\frac{p-1}{2}p+\frac{p-1}{2}}{sp+t}^2 \pmod{p^2}
\end{aligned}$$

and so

$$S_{\frac{p^2-1}{2}} \equiv (-1)^{\frac{p-1}{2}}(2^p-1)f_{\frac{p^2-1}{2}} + (-1)^{\frac{p-1}{2}}p\sum_{s=0}^{(p-1)/2} H_{2s}\sum_{t=0}^{(p-1)/2}\binom{\frac{p-1}{2}p+\frac{p-1}{2}}{sp+t}^3 \pmod{p^2}.$$

Now we assert that

$$\sum_{t=0}^{(p-1)/2}\binom{\frac{p-1}{2}p+\frac{p-1}{2}}{sp+t}^3 \equiv 0 \pmod{p^2} \quad \text{for} \quad s=0,1,2,\dots. \tag{3.4}$$

We prove the result by induction on s. From [S1] we know that the result is true for $s=0$. Suppose

$$\begin{aligned}
&\sum_{t=0}^{(p-1)/2}\binom{\frac{p-1}{2}p+\frac{p-1}{2}}{sp+t}^3\\
&\equiv \binom{\frac{p-1}{2}}{s}^3\sum_{t=0}^{(p-1)/2}\Big[\frac{p+1}{2}\binom{\frac{p-1}{2}}{t}-(\frac{p-1}{2}+s)\binom{\frac{p-1}{2}-p}{t}-s\binom{\frac{p-1}{2}-p}{t+p}\Big]^3 \equiv 0 \pmod{p^2}.
\end{aligned}$$

By Lemma 2.15,

$$\begin{aligned}
&\sum_{t=0}^{(p-1)/2}\binom{\frac{p-1}{2}p+\frac{p-1}{2}}{(s+1)p+t}^3\\
&\equiv \binom{\frac{p-1}{2}}{s+1}^3\sum_{t=0}^{(p-1)/2}\Big[\frac{p+1}{2}\binom{\frac{p-1}{2}}{t}-(\frac{p-1}{2}+s+1)\binom{\frac{p-1}{2}-p}{t}-(s+1)\binom{\frac{p-1}{2}-p}{t+p}\Big]^3\\
&= \binom{\frac{p-1}{2}}{s+1}^3\sum_{t=0}^{(p-1)/2}\Big[\frac{p+1}{2}\binom{\frac{p-1}{2}}{t}-(\frac{p-1}{2}+s)\binom{\frac{p-1}{2}-p}{t}\\
&\quad -s\binom{\frac{p-1}{2}-p}{t+p}-\Big(\binom{\frac{p-1}{2}-p}{t}+\binom{\frac{p-1}{2}-p}{t+p}\Big)\Big]^3 \pmod{p^2}.
\end{aligned}$$

Hence $\sum\limits_{t=0}^{(p-1)/2}\binom{\frac{p-1}{2}p+\frac{p-1}{2}}{(s+1)p+t}^3 \equiv 0 \pmod{p^2}$ for $s \geq \frac{p-1}{2}$. For $s < \frac{p-1}{2}$, by the inductive hypothesis we have

$$\sum_{t=0}^{(p-1)/2}\Big[\frac{p+1}{2}\binom{\frac{p-1}{2}}{t}-(\frac{p-1}{2}+s)\binom{\frac{p-1}{2}-p}{t}-s\binom{\frac{p-1}{2}-p}{t+p}\Big]^3 \equiv 0 \pmod{p^2}.$$

As $t \in \{0,1,\ldots,\frac{p-1}{2}\}$, we have $\binom{\frac{p-1}{2}}{t} \equiv \binom{\frac{p-1}{2}-p}{t} \equiv \binom{-\frac{1}{2}}{t} \pmod p$. From Lemma 2.16 we have $\binom{\frac{p-1}{2}-p}{t}+\binom{\frac{p-1}{2}-p}{t+p} = (-1)^t\Big(\binom{\frac{p-1}{2}+t}{t}-\binom{p+\frac{p-1}{2}+t}{t+p}\Big) \equiv 0 \pmod p$, and so

$$\begin{aligned}
&\sum_{t=0}^{(p-1)/2}\binom{\frac{p-1}{2}p+\frac{p-1}{2}}{(s+1)p+t}^3\\
&\equiv \binom{\frac{p-1}{2}}{s+1}^3\Big\{\sum_{t=0}^{(p-1)/2}\Big[\frac{p+1}{2}\binom{\frac{p-1}{2}}{t}-(\frac{p-1}{2}+s)\binom{\frac{p-1}{2}-p}{t}-s\binom{\frac{p-1}{2}-p}{t+p}\Big]^3\\
&\quad+3\sum_{t=0}^{(p-1)/2}\Big[\frac{p+1}{2}\binom{\frac{p-1}{2}}{t}-(\frac{p-1}{2}+s)\binom{\frac{p-1}{2}-p}{t}-s\binom{\frac{p-1}{2}-p}{t+p}\Big]\\
&\qquad\qquad\times\Big[\binom{\frac{p-1}{2}-p}{t}+\binom{\frac{p-1}{2}-p}{t+p}\Big]^2\\
&\quad-3\sum_{t=0}^{(p-1)/2}\Big[\frac{p+1}{2}\binom{\frac{p-1}{2}}{t}-(\frac{p-1}{2}+s)\binom{\frac{p-1}{2}-p}{t}-s\binom{\frac{p-1}{2}-p}{t+p}\Big]^2\\
&\qquad\qquad\times\Big[\binom{\frac{p-1}{2}-p}{t}+\binom{\frac{p-1}{2}-p}{t+p}\Big]\\
&\quad-\sum_{t=0}^{(p-1)/2}\Big[\binom{\frac{p-1}{2}-p}{t}+\binom{\frac{p-1}{2}-p}{t+p}\Big]^3\Big\}\\
&\equiv -3\binom{\frac{p-1}{2}}{s+1}^3\sum_{t=0}^{(p-1)/2}\binom{-\frac{1}{2}}{t}^2\Big[\binom{\frac{p-1}{2}-p}{t}+\binom{\frac{p-1}{2}-p}{t+p}\Big]\\
&= -3\binom{\frac{p-1}{2}}{s+1}^3\sum_{t=0}^{(p-1)/2}(-1)^t\binom{-\frac{1}{2}}{t}^2\Big[\binom{\frac{p-1}{2}+t}{t}-\binom{p+\frac{p-1}{2}+t}{t+p}\Big]\\
&= 0 \pmod{p^2}.
\end{aligned}$$

Summarizing the above proves the theorem.

4. $\{S_m\}$ is log-convex

A sequence $\{a_m\}$ $(m \geq 0)$ is called log-convex if $a_m \geq 0$ and $a_{m-1}a_{m+1} \geq a_m^2$ for $m = 1,2,3,\ldots$. In this section we show that $\{S_m\}$ and $\{P_m\}$ are log-convex sequences.

Theorem 4.1. *For $m = 2,3,4,\ldots$ we have*

$$S_m^2 < S_{m+1}S_{m-1} < \big(1+\frac{1}{m(m-1)}\big)S_m^2.$$

Proof. We first prove $S_m^2 < S_{m+1}S_{m-1}$ for $m \geq 2$. Since $S_1 = 4$, $S_2 = 20$, $S_3 = 112$, $S_4 = 676$ and $S_5 = 4304$ we see that $S_m^2 < S_{m+1}S_{m-1}$ for $m = 2, 3, 4$. From now on we assume $m \geq 5$. Suppose that $S_{m-1}^2 < S_{m-2}S_m$. By (1.3), Lemma 2.7 and the fact that $\frac{3m-4}{m-1} \geq \frac{11}{4}$ we have

$$\begin{aligned}
&S_{m+1}S_{m-1} - S_m^2 \\
&= \frac{4(3m^2+3m+1)}{(m+1)^2} S_m S_{m-1} - \frac{32m^2}{(m+1)^2} S_{m-1}^2 - S_m^2 \\
&> \Big(\frac{4(3m^2+3m+1)}{(m+1)^2} S_{m-1} - \frac{32m^2}{(m+1)^2} S_{m-2} - S_m\Big) S_m \\
&= \Big(4(3m^2+m-1)S_{m-1} + 32(1-2m^2)S_{m-2}\Big) \frac{S_m}{m^2(m+1)^2} \\
&= \Big(4(3m^2+m-1)2\sum_{k=1}^{m-1}\binom{m-2}{k-1}\binom{2k}{k}\binom{2m-2-2k}{m-1-k} \\
&\quad + 32(1-2m^2)\sum_{k=0}^{m-2}\binom{m-2}{k}\binom{2k}{k}\binom{2m-4-2k}{m-2-k}\Big)\frac{S_m}{m^2(m+1)^2} \\
&= \Big(\sum_{k=1}^{m-3}\binom{m-2}{k}\binom{2k}{k}\binom{2m-4-2k}{m-2-k}\Big(\frac{16(3m^2+m-1)(2k+1)}{k+1} + 32(1-2m^2)\Big) \\
&\quad + \Big(16(3m^2+m-1)\frac{3m-4}{m-1} + 64 - 128m^2\Big)\binom{2m-4}{m-2}\Big)\frac{S_m}{m^2(m+1)^2} \\
&> \Big(8(m^2+3m+1)\sum_{k=1}^{m-3}\binom{m-2}{k}\binom{2k}{k}\binom{2m-4-2k}{m-2-k} \\
&\quad + 4(m^2+11m+5)\binom{2m-4}{m-2}\Big)\frac{S_m}{m^2(m+1)^2} \\
&> 0.
\end{aligned}$$

Thus the inequality $S_m^2 < S_{m+1}S_{m-1}$ is proved by induction.

Next we prove the remaining inequality. It is easily seen that $\big(1+\frac{1}{m(m-1)}\big)S_m^2 - S_{m+1}S_{m-1} > 0$ for $m = 2, 3, \ldots, 13$. Now suppose $m \geq 14$ and $\big(1+\frac{1}{(m-1)(m-2)}\big)S_{m-1}^2 > S_m S_{m-2}$. By (1.3), Lemma 2.7 and the inductive hypothesis we have

$$\begin{aligned}
&\big(1+\frac{1}{m(m-1)}\big)S_m^2 - S_{m+1}S_{m-1} \\
&= \big(1+\frac{1}{m(m-1)}\big)S_m^2 - \frac{4(3m^2+3m+1)}{(m+1)^2}S_mS_{m-1} + \frac{32m^2}{(m+1)^2}S_{m-1}^2 \\
&> \Big(\frac{m^2-m+1}{m(m-1)}S_m - \frac{4(3m^2+3m+1)}{(m+1)^2}S_{m-1} + \frac{32m^2(m-1)(m-2)}{(m+1)^2(m^2-3m+3)}S_{m-2}\Big)S_m \\
&= \Big((20m^5 - 60m^4 + 52m^3 + 28m^2 - 36m + 12)S_{m-1} \\
&\quad + (-128m^5 + 320m^4 - 256m^3 - 32m^2 + 192m - 96)S_{m-2}\Big) \\
&\qquad \times \frac{S_m}{(m+1)^2(m^2-3m+3)m^3(m-1)}
\end{aligned}$$

$$= 16\sum_{k=0}^{m-2}\binom{m-2}{k}\binom{2k}{k}\binom{2m-4-2k}{m-2-k}F(m,k)\frac{S_m}{(m+1)^2(m^2-3m+3)m^3(m-1)},$$

where

$$\begin{aligned} F(m,k) = {} & (5m^5-15m^4+13m^3+7m^2-9m+3)\frac{2k+1}{k+1} \\ & -8m^5+20m^4-16m^3-2m^2+12m-6. \end{aligned}$$

For $m \geq 14$ we see that $3 < \frac{(2m-7)(2m-5)}{(m-3)(m-2)} < 4$, $5m^5-15m^4+13m^3+7m^2-9m+3>0$, $-8m^5+20m^4-16m^3-2m^2+12m-6<0$, $6m^7-75m^6+223m^5-283m^4-61m^3+427m^2-87m-42>0$, and $F(m,k+1)>F(m,k)$ for $k=0,1,\ldots,m-2$. Thus $F(m,m-3)+F(m,1)>F(m,5)+F(m,1)>0$. Since

$$\begin{aligned} F(m,k) \geq F(m,2) = {} & \frac{5}{3}(5m^5-15m^4+13m^3+7m^2-9m+3) \\ & -8m^5+20m^4-16m^3-2m^2+12m-6>0, \end{aligned}$$

from the above we derive that

$$\begin{aligned} & \big(1+\frac{1}{m(m-1)}\big)S_m^2 - S_{m+1}S_{m-1} \\ > {} & 16\Big\{\sum_{k=0}^{2}\binom{m-2}{k}\binom{2k}{k}\binom{2m-4-2k}{m-2-k}F(m,k) \\ & + \sum_{k=m-4}^{m-2}\binom{m-2}{k}\binom{2k}{k}\binom{2m-4-2k}{m-2-k}F(m,k)\Big\}\frac{S_m}{(m+1)^2(m^2-3m+3)m^3(m-1)} \\ = {} & \frac{16S_m}{(m+1)^2(m^2-3m+3)m^3(m-1)}\Big\{\binom{2m-4}{m-2}\big(F(m,m-2)+F(m,0)\big) \\ & + 2(m-2)\binom{2m-6}{m-3}\big(F(m,1)+F(m,m-3)\big) \\ & + 3(m-2)(m-3)\binom{2m-8}{m-4}\big(F(m,m-4)+F(m,2)\big)\Big\} \\ > {} & \frac{16S_m}{(m+1)^2(m^2-3m+3)m^3(m-1)} \\ & \times\Big(\big(3(m^2-5m+6)+\frac{4(2m-7)(2m-5)}{(m-3)(m-2)}\big)F(m,m-4) \\ & + \frac{4(2m-7)(2m-5)}{(m-3)(m-2)}F(m,0)\Big)\binom{2m-8}{m-4} \\ > {} & \frac{16S_m\binom{2m-8}{m-4}}{(m+1)^2(m^2-3m+3)m^3(m-1)} \\ & \times\Big(3(m^2-5m+6)F(m,m-4)+\frac{4(2m-7)(2m-5)}{(m-3)(m-2)}F(m,0)\Big) \\ = {} & \frac{S_m\binom{2m-8}{m-4}}{(m+1)^2(m^2-3m+3)m^3(m-1)}\Big(\big(6(m-2)(2m-7)+\frac{8(2m-7)(2m-5)}{(m-3)(m-2)}\big) \\ & \times(40m^5-120m^4+104m^3+56m^2-72m+24) \end{aligned}$$

$$
\begin{aligned}
&+\big(3(m-2)(m-3)+\frac{4(2m-7)(2m-5)}{(m-3)(m-2)}\big)\\
&\qquad\times(-128m^5+320m^4-256m^3-32m^2+192m-96)\Big)\\
>&\ \frac{S_m\binom{2m-8}{m-4}}{(m+1)^2(m^2-3m+3)m^3(m-1)}\\
&\times\Big(\big(6(m-2)(2m-7)+24\big)(40m^5-120m^4+104m^3+56m^2-72m+24)\\
&+\big(3(m-2)(m-3)+16\big)(-128m^5+320m^4-256m^3-32m^2+192m-96)\Big)\\
=&\ \frac{S_m\binom{2m-8}{m-4}}{(m+1)^2(m^2-3m+3)m^3(m-1)}\\
&\times(96m^7-1200m^6+3568m^5-4528m^4-976m^3+6832m^2-1392m-672)\\
>&\ 0.
\end{aligned}
$$

Hence the inequality is proved by induction.

Corollary 4.1. *Both $\{S_m\}$ and $\{P_m\}$ are log-convex.*

Proof. By (1.6), $S_m > 0$. Since $S_0 = 1$, $S_1 = 4$ and $S_2 = 20$, we see that $S_1^2 < S_0S_2$. Now applying Theorem 4.1 we see that $\{S_m\}$ is log-convex. Since $P_m = 2^mS_m$ we see that

$$P_{m-1}P_{m+1} - P_m^2 = 2^{m-1}S_{m-1} \cdot 2^{m+1}S_{m+1} - 2^{2m}S_m^2 = 2^{2m}(S_{m-1}S_{m+1} - S_m^2) \geq 0.$$

Thus $\{P_m\}$ is also log-convex.

References

[C] E. Catalan, *Sur les nombres de Segner*, Rend. Circ. Mat. Palermo **1**(1887), 190-201.

[LF1] P. Larcombe and D. French, *On the other Catalan numbers: a historical formulation re-examined*, Congr. Numer. **143**(2000), 33-64.

[LF2] P.J. Larcombe and D.R. French, *A new generating function for the Catalan-Larcombe-French sequence: proof of a result by Jovovic*, Congressus Numerantium, **166** (2004), 161-172.

[JLF] A.F. Jarvis, P.J. Larcombe and D.R. French, *On small prime divisibility of the Catalan-Larcombe-French sequence*, Indian J. Math. **47** (2005), 159-181.

[JV] F. Jarvis and H.A. Verrill, *Supercongruences for the Catalan-Larcombe-French numbers*, Ramanujan J. **22**(2010), 171-186.

[MOS] W. Magnus, F. Oberhettinger and R.P. Soni, *Formulas and Theorems for the Special Functions of Mathematical Physics* (3*rd edn.*), Springer, New York, 1966, pp.25-32.

[M] R. Meštrović, *Lucas' theorem: its generalizations, extensions and applications (1878-2014)*, arXiv:1409.3820.

[OS] R. Osburn and B. Sahu, *A supercongruence for generalized Domb numbers*, Funct. Approx. Comment. Math. **48**(2013), part 1, 29-36.

[S] B. Sagan, *Congruences via Abelian Groups*, J. Number Theory. **20**(1985), 210-237.

[S1] Z.H. Sun, *Congruences concering Legendre polynomials II*, J.Number Theory. **133** (2013), 1950-1976.

[S2] Z.H. Sun, *Super congruences involving Bernoulli and Euler polynomials*, arXiv:1407.0636.

[S3] Z.H. Sun, *Congruences involving Franel and Catalan-Larcombe-French numbers*, arXiv:1502.02499.

[Su] Z.W. Sun, *Supercongruences and Euler numbers*, Sci. China Math. **54**(2012), 2509-2535.